

兔抗 CSF1R(Phospho-Tyr708) 多克隆抗体

中文名称：兔抗 CSF1R(Phospho-Tyr708) 多克隆抗体

英文名称： Anti-CSF1R(Phospho-Tyr708) rabbit polyclonal antibody

别名： FMS; CSFR; FIM2; HDLS; C-FMS; CD115; CSF-1R; M-CSF-R

储存： 冷冻（-20℃） 避光

抗原： CSF1R(Phospho-Tyr708)

宿主： Rabbit

反应种属： Human

相关类别： 一抗

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results

	in multiple transcript variants. [provided by RefSeq, Dec 2013]
Applications:	WB
Name of antibody:	CSF1R(Phospho-Tyr708)
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 708 (K-K-Y(p)-V-R) derived from Human CSFR .
Full name:	colony stimulating factor 1 receptor
Synonyms :	FMS; CSFR; FIM2; HDLS; C-FMS; CD115; CSF-1R; M-CSF-R
SwissProt:	P07333
WB Predicted band size:	108 kDa
WB Positive control:	HepG2 cells lysates
WB Recommended dilution:	500-1000

