

兔抗 GOLGA5 多克隆抗体

中文名称：兔抗 GOLGA5 多克隆抗体

英文名称：Anti-GOLGA5 rabbit polyclonal antibody

别名：golgin A5; RFG5; GOLIM5; ret-II

抗原：GOLGA5

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

相关类别：一抗

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

The Golgi apparatus, which participates in glycosylation and transport of proteins and lipids in the secretory pathway, consists of a series of stacked cisternae (flattened membrane sacs). Interactions between the Golgi and microtubules are thought to be important for the reorganization of the Golgi after it fragments during mitosis. This gene encodes one of the golgins, a family of proteins localized to the Golgi. This protein is a coiled-coil membrane protein that has been postulated to play a role in vesicle tethering and docking. Translocations involving this gene and the ret proto-oncogene have been found in tumor tissues; the chimeric sequences have been designated RET-II and PTC5. A pseudogene of this gene is located

	ated on the short arm of chromosome 5.
Applications:	ELISA, WB, IHC
Name of antibody:	GOLGA5
Immunogen:	Synthetic peptide of human GOLGA5
Full name:	golgin A5
Synonyms:	RFG5; GOLIM5; ret-II
SwissProt:	Q8TBA6
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human thyroid cancer and Human ovarian cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	83 kDa
WB Positive control:	A549 cell lysate
WB Recommended dilution:	200-1000



