

兔抗 ABCA1 多克隆抗体

中文名称：兔抗 ABCA1 多克隆抗体

英文名称： Anti-ABCA1 rabbit polyclonal antibody

别名： ATP binding cassette subfamily A member 1; TGD; ABC1; CERP; ABC-1; HDLDT1

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： ABCA1

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ABC1 subfamily. Members of the ABC1 subfamily comprise the only major ABC subfamily found exclusively in multicellular eukaryotes. With cholesterol as its substrate, this protein functions as a cholesterol efflux pump in the cellular lipid removal pathway. Mutations in this gene have been associated with Tangier's disease and familial high-density lipoprotein deficiency.
Applications:	ELISA, WB, IHC

Name of antibody:	ABCA1
Immunogen:	Synthetic peptide of human ABCA1
Full name:	ATP binding cassette subfamily A member 1
Synonyms:	TGD; ABC1; CERP; ABC-1; HDLDT1
SwissProt:	O95477
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer
IHC Recommend dilution:	50-100
WB Predicted band size:	254 kDa
WB Positive control:	Human bladder transitional cell carcinoma grade 2-3 tissue, Human bladder transitional cell carcinoma cancer next level 2-3 tissue lysates
WB Recommended dilution:	200-1000



